

Bangladesh is today the world's most polluted country. Air pollution shortens the average Bangladeshi's life expectancy by 6.2 years, relative to what it would have been if the World Health Organization (WHO) guideline was met. Some areas of Bangladesh fare much worse than average, with air pollution shortening lives by about 7 years in the most polluted district.

KEY TAKE-AWAYS

- All of Bangladesh's 161 million people live in areas where the annual average particulate pollution level exceeds both the WHO guideline and the country's own national standard.
- Particulate pollution has sharply increased over time. Since 1998, average annual particulate pollution has increased 89 percent, cutting 3.4 years off the lives of the average Bangladesh resident over those years.
- In every one of the country's 64 districts, particulate pollution levels are at least four times the WHO guideline. The most polluted areas of the country are the divisions of Dhaka and Khulna (which include the country's first and third largest cities of the same names), where the average resident is exposed to pollution that is more than eight times the WHO guideline—cutting life expectancy by about 7 years.
- Even in the second largest city of Chittagong, where air quality is better than the national average, residents still lose 4.4 years off their lives relative to if the WHO guideline was met.

POLICY IMPACTS

The dual challenges of economic growth and environmental quality faced by Bangladesh today are no different from those once confronted by other countries during periods of industrialization. Nor is this dynamic limited to the world's wealthiest countries.

China has made tremendous progress since declaring a “war against pollution” in 2014, with cities cutting particulate pollution by about 40 percent—improving life expectancy by 2 years if the reductions persist. India, having declared its own war against pollution in January 2019, has set an ambitious target to reduce pollution by 20-30 percent. If it achieves a 25 percent reduction in pollution nationwide, it has the potential to improve life expectancy by 2 years.

Bangladesh has the opportunity to experience the same progress. If Bangladesh were to achieve the same reduction in pollution experienced by China, its residents could live 2.9 years longer; 1.8 years longer if it achieves India's target.

Figure 1 · Potential Gain in Life Expectancy through Permanently Reducing PM_{2.5} from 2018 Concentration to the WHO Guideline





Figure 2 · Average Years of Life Expectancy Lost due to PM_{2.5} Relative to the WHO Guideline, 1998-2018



PM_{2.5} Concentration and Potential Life Expectancy Gains in 25 Most Populous Districts.

Years of Life Expectancy Gain through Reducing PM_{2.5} from 2018 Concentration

Division	District	Population (Millions)	PM _{2.5} Concentration, 2018 (µg/m ³)	To WHO Guideline of 10 µg/m ³	To National Standard of 15 µg/m ³	By 25%	By 40%
Dhaka	Dhaka	13.4	83	7.2	6.7	2.0	3.3
Chittagong	Chittagong	8.4	55	4.4	3.9	1.3	2.2
Chittagong	Comilla	5.9	80	6.8	6.3	2.0	3.1
Mymensingh	Mymensingh	5.7	70	5.9	5.4	1.7	2.7
Dhaka	Tangail	4.0	77	6.6	6.1	1.9	3.0
Dhaka	Gazipur	3.9	82	7.0	6.5	2.0	3.2
Sylhet	Sylhet	3.8	58	4.7	4.2	1.4	2.3
Rajshahi	Bogra	3.8	70	5.8	5.3	1.7	2.7
Chittagong	Noakhali	3.4	73	6.2	5.7	1.8	2.9
Rajshahi	Sirajganj	3.4	77	6.6	6.1	1.9	3.0
Chittagong	Cox's Bazar	3.4	42	3.1	2.6	1.0	1.6
Rangpur	Dinajpur	3.3	67	5.6	5.1	1.6	2.6
Dhaka	Kishoreganj	3.2	74	6.2	5.7	1.8	2.9
Rangpur	Rangpur	3.2	67	5.6	5.1	1.7	2.6
Chittagong	Brahamanbari	3.2	78	6.7	6.2	1.9	3.1
Dhaka	Narayanganj	3.1	85	7.3	6.8	2.1	3.3
Khulna	Jessore	3.1	83	7.2	6.7	2.0	3.3
Rajshahi	Naogaon	2.9	70	5.9	5.4	1.7	2.8
Rajshahi	Rajshahi	2.9	75	6.3	5.8	1.8	2.9
Rajshahi	Pabna	2.8	81	6.9	6.4	2.0	3.2
Sylhet	Sunamganj	2.7	60	4.9	4.5	1.5	2.4
Chittagong	Chandpur	2.7	80	6.8	6.3	2.0	3.1
Rangpur	Gaibandha	2.6	67	5.6	5.1	1.7	2.6
Barisal	Barisal	2.6	80	6.8	6.4	2.0	3.1
Khulna	Khulna	2.6	83	7.2	6.7	2.0	3.3

“The legacy of environmental improvement in former pollution capitals is evidence that today’s pollution does not need to be tomorrow’s fate. As countries navigate the dual challenges of sustaining economic growth and protecting the environment and public health, the AQLI shows not only the damage caused by pollution but also the enormous gains that can be made with policies to address it.”

Michael Greenstone, The Milton Friedman Distinguished Service Professor in Economics, the College, and the Harris Shool; Director, EPIC

ABOUT THE AIR QUALITY LIFE INDEX (AQLI)

The AQLI is a pollution index that converts particulate air pollution into perhaps the most important metric that exists: its impact on life expectancy. Developed by the Energy Policy Institute at the University of Chicago (EPIC), the AQLI is rooted in cutting-edge, peer-reviewed research co-authored by EPIC Director Michael Greenstone that for the first time quantified the causal relationship between long-term human exposure to air pollution and life expectancy. The Index then combines this research with hyper-localized, global particulate measurements, yielding unprecedented insight into the true cost of particulate pollution in communities around the world. The Index also illustrates how air pollution policies can increase life expectancy when they meet the World Health Organization’s guideline for what is considered a safe level of exposure, existing national air quality standards, or user-defined air quality levels. This information can help to inform local communities and policymakers about the importance of air pollution policies in concrete terms.

Methodology: The life expectancy calculations made by the AQLI are based on a pair of peer-reviewed studies, Chen et al. (2013) and Ebenstein et al. (2017), co-authored by Michael Greenstone, that exploit a unique natural experiment in China. By comparing two subgroups of the population that experienced prolonged exposure to different levels of particulate air pollution, the studies were able to plausibly isolate the effect of particulates air pollution from other factors that affect health. The more recent of the two studies found that sustained exposure to an additional 10 µg/m³ of PM₁₀ reduces life expectancy by 0.64 years. In terms of PM_{2.5}, this translates to the relationship that an additional 10 µg/m³ of PM_{2.5} reduces life expectancy by 0.98 years. To learn more about the methodology used by the AQLI, visit: aqli.epic.uchicago.edu/about/methodology