



Nepal Fact Sheet

Nepal is the world's third most polluted country based on satellite-derived PM_{2.5} data. Fine particulate air pollution (PM_{2.5}) shortens the average Nepalese resident's life expectancy by 4.6 years, relative to what it would be if the World Health Organization (WHO) guideline of 5 µg/m³ was met.¹ Some areas of Nepal fare much worse than average, with air pollution shortening lives by 6.8 years in the nine districts with the highest concentration of particulate pollution.² These districts lie in southern Nepal and share their borders with the highly-polluted Northern Plains of India.³

KEY TAKEAWAYS

- All of Nepal's 30.5 million people live in areas where the annual average particulate pollution level exceeds the WHO guideline. Despite a high health burden due to particulate pollution, the country does not have a national standard for annual average PM_{2.5}.
- Measured in terms of life expectancy, particulate pollution is the greatest threat to human health in Nepal, taking 4.6 years off the life of the average Nepalese resident. In contrast, tobacco use reduces life expectancy by 2.8 years, while high blood pressure reduces life expectancy by 1.7 years.
- Particulate pollution has increased over time. From 1998 to 2021, average annual particulate pollution increased by 75.2 percent, further reducing life expectancy by 2.2 years.
- If Nepal were to reduce particulate pollution to meet the WHO guideline, residents in the mid and eastern Terai region—where nearly 40 percent of Nepal's population resides—would gain 6.5 years of life expectancy.⁴ In the capital city of Kathmandu—Nepal's most populous city—residents would gain 3.5 years of life expectancy.

Figure 1 · Potential gain in life expectancy from permanently reducing PM_{2.5} from 2021 concentration to the WHO guideline





Figure 2 · Potential gain in life expectancy from reducing PM_{2.5} from 2021 levels to the WHO guideline in all provinces of Nepal




1 This data is based on the AQLI 2021 dataset. All annual average PM_{2.5} values (measured in micrograms per cubic meter: µg/m³) are population weighted.
 2 We include the districts of Kapilbastu, Rupandehi, Nawalparasi West, Parsa, Bara, Rautahat, Sarlahi, Mahottari, Siraha, Saptari to calculate this average.
 3 We define the Indo-Gangetic plain region as the following seven states and union territories: Bihar, Chandigarh, Delhi, Haryana, Punjab, Uttar Pradesh, and West Bengal.
 4 We define the mid and eastern Terai region as the following districts: Kapilbastu, Rupandehi, Nawalparasi West, Nawalparasi East, Parsa, Bara, Rautahat, Sarlahi, Mahottari, Dhanusha, Siraha, Saptari, Sunsari, Morang, Jhapa.

Potential life expectancy impacts of particulate pollution reductions in the 25 most populous regions of Nepal

District	Population (millions)	Annual average 2021 PM _{2.5} concentration (µg/m ³)	Life expectancy gains from reducing PM _{2.5} from 2021 concentration to WHO PM _{2.5} guideline of 5 µg/m ³ (years)	Life expectancy gains from reducing PM _{2.5} from 2021 concentration by 30 percent (years)
Kathmandu	2	40.5	3.5	1.2
Morang	1.1	61.3	5.5	1.8
Rupandehi	1	68.4	6.2	2
Jhapa	0.9	58.7	5.3	1.7
Kailali	0.9	54.9	4.9	1.6
Sarlahi	0.9	78	7.2	2.3
Dhanusha	0.9	80	7.4	2.4
Sunsari	0.9	61.4	5.5	1.8
Bara	0.8	78.7	7.2	2.3
Rautahat	0.8	80.8	7.4	2.4
Saptari	0.7	68	6.2	2
Siraha	0.7	73.5	6.7	2.2
Mahottari	0.7	81.7	7.5	2.4


District	Population (millions)	Annual average 2021 PM _{2.5} concentration (µg/m ³)	Life expectancy gains from reducing PM _{2.5} from 2021 concentration to WHO PM _{2.5} guideline of 5 µg/m ³ (years)	Life expectancy gains from reducing PM _{2.5} from 2021 concentration by 30 percent (years)
Parsa	0.7	78.1	7.2	2.3
Chitawan	0.7	55.4	4.9	1.6
Kapilbastu	0.7	67.6	6.1	2
Dang	0.6	44.4	3.9	1.3
Banke	0.6	61.8	5.6	1.8
Kaski	0.6	34.1	2.9	1
Kanchanpur	0.5	52.7	4.7	1.5
Lalitpur	0.5	38.7	3.3	1.1
Bardiya	0.5	60.2	5.4	1.8
Makawanpur	0.5	47.8	4.2	1.4
Kabhrepalanchok	0.4	37.9	3.2	1.1
Surkhet	0.4	40.1	3.4	1.2

Figure 3 · Top 5 threats to life expectancy in Nepal



Sources: Global Burden of Disease (<https://vizhub.healthdata.org/gbd-results/>) level-2 causes and risks data and WHO Life Tables (<https://apps.who.int/gho/data/node.main.LIFECOUNTRY?lang=en>) were combined with the Life table method to arrive at these results. *PM_{2.5} relative to WHO Guideline' bar displays the reduction in life expectancy relative to the WHO guideline as calculated by latest AQLI (2021) data.

Figure 4 · Annual average PM_{2.5} concentration in Nepal, 1998-2021



ABOUT THE AIR QUALITY LIFE INDEX (AQLI)

The AQLI is a pollution index that translates particulate air pollution into perhaps the most important metric that exists: its impact on life expectancy. Developed by the University of Chicago's Milton Friedman Distinguished Service Professor in Economics Michael Greenstone and his team at the Energy Policy Institute at the University of Chicago (EPIC), the AQLI is rooted in research that quantifies the causal relationship between long-term human exposure to air pollution and life expectancy. The Index then combines this research with hyper-localized, satellite measurements of global particulate matter (PM_{2.5}), yielding unprecedented insight into the true cost of pollution in communities around the world. The Index also illustrates how air pollution policies can increase life expectancy when they meet the World Health Organization's guideline for what is considered a safe level of exposure, existing national air quality standards, or user-defined air quality levels. This information can help to inform local communities and policymakers about the importance of air pollution policies in concrete terms.

Methodology: The life expectancy calculations made by the AQLI are based on a pair of peer-reviewed studies, Chen et al. (2013) and Ebenstein et al. (2017), co-authored by Michael Greenstone, that exploit a unique natural experiment in China. By comparing two subgroups of the population that experienced prolonged exposure to different levels of particulate air pollution, the studies were able to plausibly isolate the effect of particulate air pollution from other factors that affect health. Ebenstein et al. (2017) found that sustained exposure to an additional 10 µg/m³ of PM₁₀ reduces life expectancy by 0.64 years. In terms of PM_{2.5}, this translates to the relationship that an additional 10 µg/m³ of PM_{2.5} reduces life expectancy by 0.98 years. This metric is then combined with sea-salt and mineral dust removed satellite-derived PM_{2.5} data. All 2021 annual average PM_{2.5} values are population-weighted and AQLI's source of population data is <https://landscan.ornl.gov/>. We are grateful to the Atmospheric Composition Analysis Group, based at the Washington University in St. Louis for providing us with the satellite data. The original dataset can be found here: <https://sites.wustl.edu/acag/datasets/surface-pm2-5/>. To learn more deeply about the methodology used by the AQLI, visit: aqli.epic.uchicago.edu/about/methodology.